


POTENTIAL IMPACTS OF GLOBAL CLIMATE CHANGE ON THE WATER RESOURCES OF PUERTO RICO




Ferdinand Quiñones, PE

Objetives


- Briefly review current issues on Global Climate Change
- Describe the simplified Thornthwaite climatic index used to evaluate potential climate changes in Puerto Rico
- Present perspective of the water resources of Puerto Rico in the climate change environment
- Results of the evaluations and conclusions
- Questions and Answers

HISTORIC AND RECENT TEMPERATURE DATA




HISTORIC CO2 CONCENTRATIONS IN THE ATMOSPHERE

CO2 concentrations 647,000 BC to 2006 AD
Antarctic temperature 421,000 BC to 2000 AD*



* Antarctic temperature is measured as the change from average conditions for the period 1850 AD - 2000 AD

TYPICAL GRIDS OF CLIMATE MODELS



**SIMPLIFIED EMPIRICAL MODEL
TO ESTIMATE THE POTENTIAL EFFECTS
OF CLIMATE CHANGE ON THE LIFE ZONES
OF PUERTO RICO**

THORNTHWAITE PRECIPITATION- EVAPORATION INDEX (IPET)

$$\text{IPET} = 10 \sum P/E, \text{ from } n=1,12$$

where P – Average monthly Precipitation, in inches
E – Average monthly Evaporation, in inches

Since values of Evaporation are not readily available, Thornthwaite developed an equation to estimate the IPET index using Precipitation (P) and Temperature values (T), readily available from NWS:


$$\text{IPET} = 115 \sum [(P)/(T-10)], \text{ from } n=1,12$$

THORNTHWAITE IPET INDEX APPLIED TO PUERTO RICO


Natural life zones for Puerto Rico based on Holdridge Classification System, which is correlated to Thornthwaite IPET values:

LIFE ZONE	VEGETATION CHARACTERISTICS	IPET VALUE
A (WET)	RAINFOREST	>127
C (HUMID)	SUB-TROPICAL RAINFOREST	64-127
D (SUB-HUMID)	HUMID FOREST	32-63
E (SEMI-ARID)	SUBTROPICAL FOREST	16-31
F (ARID)	DRY SUBTROPICAL FOREST	<16


LIFE ZONES IN PUERTO RICO




AVERAGE PRECIPITATION IN PUERTO RICO, 1971-2000




COMPOSITE OF PRECIPITATION AND LIFE ZONES IN PUERTO RICO



AVERAGE TEMPERATURES IN PUERTO RICO




CORRELATIONS USED TO CALCULATE THE PE INDEX




Modificado de Giusti, 1978.


AVERAGE EVAPOTRANSPIRATION IN PUERTO RICO




GRID USED IN CALCULATIONS OF PE INDEX




POTENTIAL CHANGES IN THE LIFE ZONES OF PUERTO RICO DUE TO TEMPERATURE INCREASES




WATER USE IN PUERTO RICO 1960-2005



Surface and Ground Water Use in Puerto Rico



BASINS AND STREAMS IN PUERTO RICO




TOTAL ANNUAL RUNOFF IN PUERTO RICO




RUNOFF, MINIMUM FLOWS AND WITHDRAWALS OF WATER IN PUERTO RICO

RUNOFF, MINIMUM FLOWS AND WITHDRAWALS,
ACRE-FEET PER YEAR




TOTAL ANNUAL RUNOFF IN PUERTO RICO




**POTENTIAL IMPACTS
OF CLIMATE CHANGE
ON THE
GROUND WATER RESOURCES
OF PUERTO RICO**


AQUIFERS IN PUERTO RICO




SALINITY INTRUSION IN COASTAL AQUIFERS



SALINITY INTRUSION IN PUERTO RICO



SALINE INTRUSION AND IMPACT TO THE COASTAL AQUIFERS OF PUERTO RICO




GROUND WATER LEVELS IN THE COASTAL AQUIFER OF JUANA DIAZ, SOUTHERN PUERTO RICO





POTENTIOMETRIC SURFACE OF THE ALLUVIAL AQUIFER AND HYDROLOGIC CONDITIONS IN THE JUANA DÍAZ AREA, PUERTO RICO, JUNE 29 - JULY 1, 2005

By José M. Rodríguez, Luis Santiago-Rivera, and Fernando Gómez-Gómez




ELEVATION OF THE WATER TABLE FROM THE SHORELINE







Base modified from U.S. Geological Survey digital data




VERTICAL DISPLACEMENT OF SALINE INTERFACE WITH 5 FEET OF SEA LEVEL RISE




EFFECT OF SEA LEVEL RISE OF 5 FEET ON THE SALINE WATER INTERPHASE ON THE AQUIFERS OF PUERTO RICO



SALINITY INTRUSION INTO THE RIO GRANDE DE MANATI NEAR MANATI



SALINITY WEDGE IN THE MANATI RIVER AND PROJECTED CHANGE (PURPLE) WITH A 5-FOOT INCREASE IN SEA LEVEL




Conclusions

- Increases in temperature and reduced precipitation will induce moderate changes in the natural life zones of Puerto Rico, with the more severe effects on the dry-forest areas in the southwest region
- Potential climate change effects on the Island's water resources are projected to be minimal due to abundance of water
- Ground water resources could be impacted by salinity intrusion induced by raising sea levels.

ANY WATERY QUESTIONS?????



POTENTIAL IMPACTS OF GLOBAL CLIMATE CHANGE ON THE WATER RESOURCES OF PUERTO RICO




Ferdinand Quiñones, PE

Objetives


- Briefly review current issues on Global Climate Change
- Describe the simplified Thornthwaite climatic index used to evaluate potential climate changes in Puerto Rico
- Present perspective of the water resources of Puerto Rico in the climate change environment
- Results of the evaluations and conclusions
- Questions and Answers

HISTORIC AND RECENT TEMPERATURE DATA




HISTORIC CO₂ CONCENTRATIONS IN THE ATMOSPHERE

CO₂ concentrations 647,000 BC to 2006 AD
Antarctic temperature 421,000 BC to 2000 AD*



* Antarctic temperature is measured as the change from average conditions for the period 1850 AD - 2000 AD

TYPICAL GRIDS OF CLIMATE MODELS



**SIMPLIFIED EMPIRICAL MODEL
TO ESTIMATE THE POTENTIAL EFFECTS
OF CLIMATE CHANGE ON THE LIFE ZONES
OF PUERTO RICO**

THORNTHWAITE PRECIPITATION- EVAPORATION INDEX (IPET)

$$\text{IPET} = 10 \sum P/E, \text{ from } n=1,12$$

where P – Average monthly Precipitation, in inches
E – Average monthly Evaporation, in inches

Since values of Evaporation are not readily available, Thornthwaite developed an equation to estimate the IPET index using Precipitation (P) and Temperature values (T), readily available from NWS:


$$\text{IPET} = 115 \sum [(P)/(T-10)], \text{ from } n=1,12$$

THORNTHWAITE IPET INDEX APPLIED TO PUERTO RICO


Natural life zones for Puerto Rico based on Holdridge Classification System, which is correlated to Thornthwaite IPET values:

LIFE ZONE	VEGETATION CHARACTERISTICS	IPET VALUE
A (WET)	RAINFOREST	>127
C (HUMID)	SUB-TROPICAL RAINFOREST	64-127
D (SUB-HUMID)	HUMID FOREST	32-63
E (SEMI-ARID)	SUBTROPICAL FOREST	16-31
F (ARID)	DRY SUBTROPICAL FOREST	<16


LIFE ZONES IN PUERTO RICO




AVERAGE PRECIPITATION IN PUERTO RICO, 1971-2000




COMPOSITE OF PRECIPITATION AND LIFE ZONES IN PUERTO RICO



AVERAGE TEMPERATURES IN PUERTO RICO




CORRELATIONS USED TO CALCULATE THE PE INDEX




Modificado de Giusti, 1978.


AVERAGE EVAPOTRANSPIRATION IN PUERTO RICO




GRID USED IN CALCULATIONS OF PE INDEX




POTENTIAL CHANGES IN THE LIFE ZONES OF PUERTO RICO DUE TO TEMPERATURE INCREASES




WATER USE IN PUERTO RICO 1960-2005



Surface and Ground Water Use in Puerto Rico



BASINS AND STREAMS IN PUERTO RICO




TOTAL ANNUAL RUNOFF IN PUERTO RICO




RUNOFF, MINIMUM FLOWS AND WITHDRAWALS OF WATER IN PUERTO RICO

RUNOFF, MINIMUM FLOWS AND WITHDRAWALS,
ACRE-FEET PER YEAR




TOTAL ANNUAL RUNOFF IN PUERTO RICO




**POTENTIAL IMPACTS
OF CLIMATE CHANGE
ON THE
GROUND WATER RESOURCES
OF PUERTO RICO**


AQUIFERS IN PUERTO RICO




SALINITY INTRUSION IN COASTAL AQUIFERS



SALINITY INTRUSION IN PUERTO RICO



SALINE INTRUSION AND IMPACT TO THE COASTAL AQUIFERS OF PUERTO RICO




GROUND WATER LEVELS IN THE COASTAL AQUIFER OF JUANA DIAZ, SOUTHERN PUERTO RICO





POTENTIOMETRIC SURFACE OF THE ALLUVIAL AQUIFER AND HYDROLOGIC CONDITIONS IN THE JUANA DÍAZ AREA, PUERTO RICO, JUNE 29 - JULY 1, 2005

By José M. Rodríguez, Luis Santiago-Rivera, and Fernando Gómez-Gómez




ELEVATION OF THE WATER TABLE FROM THE SHORELINE







Base modified from U.S. Geological Survey digital data.




VERTICAL DISPLACEMENT OF SALINE INTERFACE WITH 5 FEET OF SEA LEVEL RISE




EFFECT OF SEA LEVEL RISE OF 5 FEET ON THE SALINE WATER INTERPHASE ON THE AQUIFERS OF PUERTO RICO



SALINITY INTRUSION INTO THE RIO GRANDE DE MANATI NEAR MANATI



SALINITY WEDGE IN THE MANATI RIVER AND PROJECTED CHANGE (PURPLE) WITH A 5-FOOT INCREASE IN SEA LEVEL



Conclusions

- Increases in temperature and reduced precipitation will induce moderate changes in the natural life zones of Puerto Rico, with the more severe effects on the dry-forest areas in the southwest region
- Potential climate change effects on the Island's water resources are projected to be minimal due to abundance of water
- Ground water resources could be impacted by salinity intrusion induced by raising sea levels.

ANY WATERY QUESTIONS?????

